

## From chiral quarks to high-energy processes

Wojciech Broniowski

Institute of Nuclear Physics PAN, Cracow & Jan Kochanowski  
University, Kielce

with **Enrique Ruiz Arriola** and **Alexander E. Dorokhov**


Excited QCD, Zakopane, 8-14 February 2009

- *Gravitational and higher-order form factors of the pion in chiral quark models*, WB, **Enrique Ruiz Arriola**, Phys. Rev. D78 (2008) 094011
- *Generalized parton distributions of the pion in chiral quark models and their QCD evolution*, WB, ERA, **Krzysztof Golec-Biernat**, Phys. Rev. D77 (2008) 034023
- *Pion-photon Transition Distribution Amplitudes in the Spectral Quark Model*, WB, ERA, Phys. Lett. B649 (2007) 49
- *Photon distribution amplitudes and light-cone wave functions in chiral quark models*, **Alexander E. Dorokhov**, WB, ERA, Phys. Rev. D74 (2006) 054023

Other groups:


- Praszalowicz, Rostworowski, Bzdak, Kotko (Jagellonian)
- Noguera, Vento, Theussl, Courtoy (Valencia)
- Tiburzi, Miller (Seattle)
- Bochum, Tübingen groups (nucleon)

# Chiral quark models



- soft regime  $\rightarrow$  chiral sym. breaking
- NJL (Nobel 2008), instanton liquid, DSE
- relatively few parameters (traded for  $f_\pi, m_\pi, \dots$ )
- very many processes can be computed!
- no confinement - careful not to open the  $q\bar{q}$  threshold

# Example: Deep Inelastic Scattering



$$Q^2 = -q^2, \quad x = \frac{Q^2}{2p \cdot q}, \quad Q^2 \rightarrow \infty$$


Factorization of soft and hard processes, Wilson's OPE, twist expansion

$$\langle J(q)J(-q) \rangle = \sum_i C_i(Q^2; \mu) \langle \mathcal{O}_i(\mu) \rangle, \quad F(x, Q) = F_0(x, \alpha(Q)) + \frac{F_2(x, \alpha(Q))}{Q^2} + \dots$$

The soft matrix element can be computed in low-energy models!

$$F_i(x, \alpha(Q_0))|_{\text{model}} = F_i(x, \alpha(Q_0))|_{\text{QCD}}, \quad Q_0 - \text{the matching scale}$$

# QCD evolution




inclusion of gluons

- Here: DGLAP (good for intermediate  $x$ )
- Chiral quark models provide **dynamically** the non-perturbative **initial conditions** for the QCD evolution
- Inclusive and exclusive **high-energy** processes and **lattice calculations** provide the relevant data to verify the scheme

# Exclusive processes in QCD

(see earlier talks by Machado and Wagner)



Deeply  
 Virtual  
 Compton  
 Scattering

Hard  
 Meson  
 Production

## Definition of Generalized Parton Distributions

Twist-2 even-parity GPDs of the pion

non-singlet:

$$\mathcal{H}^{q,I=1}(x, \zeta, t) = \int \frac{dz^-}{4\pi} e^{ixp^+ z^-} \langle \pi^+(p+q) | \bar{\psi}(0) [0, z] \gamma^+ \tau_3 \psi(z) | \pi^+(p) \rangle \Big|_{z^+=0, z^\perp=0}$$

(similarly for singlet quarks and gluons)

$$p^2 = m_\pi^2, \quad q^2 = -2p \cdot q = t, \quad \zeta = q^+ / p^+$$

$\zeta$  - momentum transfer along the light cone

( $[0, z] = 1$  in the light-cone gauge)

## Reviews:

- K. Goeke, M. V. Polyakov, and M. Vanderhaeghen, Prog. Part. Nucl. Phys. 47 (2001) 401, hep-ph/0106012
- M. Diehl, Phys. Rept. 388 (2003) 41, hep-ph/0307382
- A. V. Belitsky, A. V. Radushkin, Phys.Rept.418(2005)1, hep-ph/0504030

GPDs provide very rich information of the structure of hadrons, encoding form factors, PDFs, ... Data may come from such processes as  $ep \rightarrow ep\gamma$ ,  $\gamma p \rightarrow pl^+l^-$ ,  $ep \rightarrow epl^+l^-$ , or from **lattices**. Small cross sections of exclusive processes require very high accuracy experiments. First results for the **nucleon** are coming from HERMES and CLAS, also COMPASS, H1, ZEUS



## Formal features

*Symmetric* notation:  $\xi = \frac{\zeta}{2-\zeta}$ ,  $X = \frac{x-\zeta/2}{1-\zeta/2}$ , with  $0 \leq \xi \leq 1$ ,  $-1 \leq X \leq 1$

$$H^{I=0}(X, \xi, t) = -H^{I=0}(-X, \xi, t), \quad H^{I=1}(X, \xi, t) = H^{I=1}(-X, \xi, t).$$

For  $X \geq 0$  we have  $\mathcal{H}^{I=0,1}(X, 0, 0) = q(X)$  - the usual PDF

The following **sum rules** hold:

$$\forall \xi : \quad \int_{-1}^1 dX H^{I=1}(X, \xi, t) = 2F_V(t),$$

$$\int_{-1}^1 dX X H^{I=0}(X, \xi, t) = 2\theta_2(t) - 2\xi^2\theta_1(t),$$

where  $F_V(t)$  is the **electromagnetic form factor**, while  $\theta_1(t)$  and  $\theta_2(t)$  are the **gravitational form factors** (related to the charge conservation and the momentum sum rule in DIS)

The **polynomiality** conditions (Lorentz invariance, time reversal, and hermiticity):

$$\int_{-1}^1 dX X^{2j} H^{I=1}(X, \xi, t) = 2 \sum_{i=0}^j A_{2j+1, 2i}(t) \xi^{2i},$$

(similarly for singlet)

$A$ 's – **generalized form factors (GFFs)**

Another way to look at GFFs:

$$\begin{aligned} \langle \pi^+(p') | \bar{u}(0) \gamma^{\{\mu} i \overleftrightarrow{D}^{\mu_1} i \overleftrightarrow{D}^{\mu_2} \dots i \overleftrightarrow{D}^{\mu_{n-1}} \} u(0) | \pi^+(p) \rangle = \\ 2P^{\{\mu} P^{\mu_1} \dots P^{\mu_{n-1}} \} A_{n0}(t) + 2 \sum_{\substack{k=2 \\ \text{even}}}^n q^{\{\mu} q^{\mu_1} \dots q^{\mu_{k-1}} P^{\mu_k} \dots P^{\mu_{n-1}} \} 2^{-k} A_{nk}(t) \end{aligned}$$

GPDs may be viewed as an infinite collection of GFFs

The **positivity bound**:

$$|H_q(X, \xi, t)| \leq \sqrt{q(x_{\text{in}})q(x_{\text{out}})}, \quad \xi \leq X \leq 1.$$


where  $x_{\text{in}} = (x + \xi)/(1 + \xi)$ ,  $x_{\text{out}} = (x - \xi)/(1 - \xi)$ .

Finally, a **low-energy theorem**  $H_{I=1}(2z - 1, 1, 0) = \phi(z)$  holds, where  $\phi$  is the pion **distribution amplitude (DA)**


Above relations and bounds impose severe constraints on the form of the GPDs

**All are satisfied in our quark-model calculation**

# QM evaluation of the GPDs



Large- $N_c =$  one loop



Direct (a), crossed (b), and contact (c) contribution ( $D$ -term) to the GPD of the pion (wavy line:  $\gamma^+$ )

## PDF, QM

With  $\zeta = t = 0$ , the GPD becomes the PDF. The Nambu–Jona-Lasinio model (Davidson, Arriola, 1995) gives


$$q(x) = 1$$

## PDF, QM

With  $\zeta = t = 0$ , the GPD becomes the PDF. The Nambu–Jona-Lasinio model (Davidson, Arriola, 1995) gives


$$q(x) = 1$$

LO DGLAP QCD evolution (good at intermediate  $x$ ) of the non-singlet part to growing scales



## PDF, QM vs. E615

LO DGLAP QCD evolution of the non-singlet part to the scale  $Q^2 = (4 \text{ GeV})^2$  of the E615 Fermilab experiment:



points: Drell-Yan from E615  
dashed: reanalysis of data  
[Wijesooriya et al., 2005]  
band: valence QM PDF evolved to  
 $Q = 4 \text{ GeV}$  from the QM scale  
 $Q_0 = 313_{-10}^{+20} \text{ MeV}$

## The quark-model scale $Q_0$

Various ways to fix: PDF, DA, moments

From experiment, the momentum fraction carried by the valence quarks is [SMRS 1992, GRS 1999]

$$\langle x \rangle_v = 0.47(2) \quad \text{at} \quad Q^2 = 4 \text{ GeV}^2$$


QM scale = no gluons, may evolve backwards until  $\langle x \rangle_v = 1$   
 → quark-model scale for NJL

$$Q_0 = 313_{-10}^{+20} \text{ MeV}$$

(here for the so called local model, for other QM  $Q_0$  may vary)  
 At this scale  $\alpha(Q_0^2)/(2\pi) = 0.34$ , which makes the evolution very fast for the scales close to the initial value – calls for improvement!




## PDF, QM vs. lattice



points: transverse lattice  
[Dalley, van de Sande, 2003]  
yellow: QM evolved to 0.35 GeV  
pink: QM evolved to 0.5 GeV  
dashed: GRS parameterization at  
0.5 GeV

# Pion Distribution Amplitude

[Bakulev, Mikhailov, Stefanis, ...]




Definition (for  $\pi^+$ , leading twist):

$$\langle 0 | \bar{d}(z) \gamma_\mu \gamma_5 u(-z) | \pi^+(q) \rangle = i\sqrt{2} f_\pi(q^2) q_\mu \int_0^1 dx e^{i(2x-1)q \cdot z} \phi(x)$$

Normalization  $\int_0^1 dx \phi(x) = 1$ , since  $\langle 0 | A_\mu^-(0) | \pi^+(q) \rangle = i f_\pi(q^2) q_\mu$   
 PDA is also relevant for the  $\pi^0 \gamma \gamma^*$  transition form factor measured by CLEO and CELLO


Similar studies in [Praszałowicz, Rostworowski, 2003]

## PDA, QM vs. E791 and lattice data




points: E791 data from di-jet  
production in  $\pi + A$   
band: QM at  $Q = 2$  GeV  
dashed line: asymptotic form  
( $Q \rightarrow \infty$ )

# PDA, QM vs. E791 and lattice data



points: E791 data from di-jet  
 production in  $\pi + A$   
 band: QM at  $Q = 2$  GeV  
 dashed line: asymptotic form  
 ( $Q \rightarrow \infty$ )



points: transverse lattice data  
 [Dalley, van de Sande, 2003]  
 band: QM at  $Q = 0.5$  GeV

## GPD in chiral quark models

Analytic formulas derived, **no factorization of the  $t$ -dependence**  
 - sheds light on possible parameterizations.

Building block of the GPD in Spectral Quark Model (SQM):

$$J_{\text{SQM}}(x, \zeta; t) = (\theta[x(\zeta - x)]\chi_1 + \theta[(1 - x)(x - \zeta)]\chi_2)$$

$$\chi_2 = \frac{2(x - 1) [3(\zeta - 1)M_V^2 + t(x - 1)^2]}{[(\zeta - 1)M_V^2 + t(x - 1)^2]^2},$$







$$\chi_1 = \frac{(x(\zeta - 2) + \zeta) (3M_V^2(\zeta - 1)\zeta^2 + t((\zeta^2 + 8\zeta - 8)x^2 + 2(4 - 5\zeta)\zeta x + \zeta^2))}{((\zeta - 1)M_V^2 + t(x - 1)^2)^2 \left(\zeta^2 + \frac{4tx(x - \zeta)}{M_V^2}\right)^{3/2}} + \frac{1}{2}\chi_2$$

$M_v$  – mass of the  $\rho$  meson

Similar studies in [Praszałowicz, Rostworowski, 2003] in a non-local model

**Next slide:**

LO DGLAP-ERBL evolution for SQM with  $\xi = 1/3$ . Solid - initial condition, dashed - evolved to  $Q^2 = (4\text{GeV})^2$ , dotted - asymptotic form. Code: [Golec-Biernat, Martin, 1999]



## Gravitational form factors

Electromagnetic current:

$$J_V^\mu = \sum_{q=u,d,\dots} \bar{q}(x) \frac{\tau_a}{2} \gamma^\mu q(x)$$

Energy-momentum tensor:

$$\Theta^{\mu\nu} = \sum_{q=u,d,\dots} \bar{q}(x) \frac{i}{2} (\gamma^\mu \partial^\nu + \gamma^\nu \partial^\mu) q(x) + \text{gluons}$$

Two structures (form factors):


$$\langle \pi^b(p') | \Theta^{\mu\nu}(0) | \pi^a(p) \rangle = \frac{1}{2} \delta^{ab} [(g^{\mu\nu} q^2 - q^\mu q^\nu) \Theta_1(q^2) + 4P^\mu P^\nu \Theta_2(q^2)]$$

traceless tensor –  $\Theta_1$  and scalar –  $\Theta_2$

Lattice, exclusive processes



## Full-QCD Euclidean lattice results




The EM FF (left) and the quark part of the gravitational form factor  $\Theta_1$  (right) in SQM (solid line) and NJL (dashed line), compared to data from [Brömmel et al., 2005-7]

Quark-model relation:  $\langle r^2 \rangle_{\Theta} = \frac{1}{2} \langle r^2 \rangle_V$

Matter more concentrated than charge!

## Higher-order form factors - predictions



The quark GFFs  $A_{3,2i}$  and  $A_{4,2i}$  at the quark-model scale  $Q_0 \sim 320$  MeV (a) and at the lattice scale  $Q = 2$  GeV (b)

## Quark moments at $t = \xi = 0$

With the notation  $\langle x^n \rangle = A_{n+1,0}(0)$ , one finds at the lattice scale of  $Q = 2$  GeV [Brömmel et al., 2007]

$$\begin{aligned}\langle x \rangle &= 0.271 \pm 0.016 \\ \langle x^2 \rangle &= 0.128 \pm 0.018 \\ \langle x^3 \rangle &= 0.074 \pm 0.027\end{aligned}$$

(lattice)

while in QM after the LO DGLAP evolution to the lattice scale

$$\begin{aligned}\langle x \rangle &= 0.28 \pm 0.02 \\ \langle x^2 \rangle &= 0.10 \pm 0.02 \\ \langle x^3 \rangle &= 0.06 \pm 0.01\end{aligned}$$

(chiral quark models)

Agreement within uncertainties

## Other quantities

- Photon DAs (with A. E. Dorokhov)
- Transition Distribution Amplitudes (TDA) [Pire, Szymanowski, 2005] (as the GPD, but between the  $\pi$  and  $\gamma$  states)
- $b$ -representation of GPDs and transverse lattices

- 1 Link between high- and low-energy analyses
- 2 Quark models provide (reasonable) initial conditions for the QCD evolution
- 3 Analytic formulas – useful for general properties, (e.g., no factorization of the  $t$ -dependence)
- 4  $Q_0$  low – need for improvement of the evolution, non-local models have higher  $Q_0$
- 5 With naive DGLAP-ERBL evolution the overall agreement with the data and lattice simulations **very reasonable** (PDF, DA, GFFs, GPD, photon DA, TDA, ...)
- 6 In QM the mean squared EM radius is twice the gravitational one
- 7 Predictions can be further tested with future lattice simulations for higher-order form factors. The behavior is non-trivial, with form factors having different signs, magnitude, and asymptotic fall-off.
- 8 GPDs of the **nucleon**: more challenging (Bochum, Tübingen – soliton) but experimental data exist