


High density phase of QCD in the PNJL model

How does deconfinement occurs at high density

Hubert Hansen <hansen@ipnl.in2p3.fr>



Institut de Physique
Nucléaire de Lyon,
CNRS/IN2P3 and université
Claude Bernard de Lyon
Collaborators: G. Goessens,
A. Gatti, P. Costa, M. Ruivo,
C. Da Sousa

Motivation: phase transition in RHIC at finite T and high density

Confinement and asymptotic freedom are difficult questions analytically and in simulation (Lattice QCD).

Besides finite density increases the difficulty (no lattice result at large density).

Surprisingly, the PNJL model is “easy” to treat, understand and it gives a good phenomenological picture where it can be compared with LQCD, even if it only implements a statistical confinement.

⇒ what kind of mechanism can we describe concerning the link between chiral transition and deconfinement, in particular at high density.

- QCD at high density in the non-perturbative sector ⇒ one needs effective model.
- Phase transition study: need chiral symmetry + confinement (associated to \mathbb{Z}_3 symmetry) ⇒ we use the PNJL model, because we want to study the interplay between chiral restoration and deconfinement.
- Only with two flavors: indeed we want to detailed the mechanisms occurring at high density (driven by the Fermi momentum) so we do not want to add other effects yet (strangeness, vector, etc.) that could hide those.
- We will study mean field predictions and then use mesonic correlations as a probe of the phase diagram.

The center symmetry of QCD at finite temperature

Finite temperature T : Equilibrium canonical ensemble $\implies x_0 \rightarrow ix_4$ (imaginary time or euclidean metric: quantum fluctuations \leftrightarrow thermal bath), $x_4 \in [0, \beta]$; $\beta = 1/T$

The Euclidean QCD Lagrangian is invariant under a gauge transformation h :

$A_\mu \longrightarrow {}^h A_\mu = h A_\mu h^\dagger - ih \partial_\mu h^\dagger$ and $q \longrightarrow {}^h q = h q$ with A_μ a gauge field, q a quark field.

From boundary condition in imaginary time:

$$A(\vec{x}, x_4 + \beta) = A(\vec{x}, x_4) \text{ and } q(\vec{x}, x_4 + \beta) = -q(\vec{x}, x_4).$$

Hence the constraints:

$${}^h A(\vec{x}, x_4 + \beta) = {}^h A(\vec{x}, x_4) \text{ and } {}^h q(\vec{x}, x_4 + \beta) = - {}^h q(\vec{x}, x_4).$$

Searching a solution of the constraints equations with

$$h(\vec{x}, x_4 + \beta) = f h(\vec{x}, x_4)$$

where $f \in$ center of $SU_c(3) \equiv \mathbb{Z}_3 = \{z_n \mathbb{1}, n = 1, 2, 3\}$ ($z_n = e^{2in\pi/3}$).

Since $[f, A_\mu] = 0$ by definition, it follows:

$${}^h A(\vec{x}, x_4 + \beta) = {}^h A(\vec{x}, x_4) \text{ (satisfies the constraint)}$$

$$\text{and } {}^h q(\vec{x}, x_4 + \beta) = -z {}^h q(\vec{x}, x_4) \text{ (breaking of } \mathbb{Z}_3).$$

\mathbb{Z}_3 is explicitly broken in the presence of light (dynamic) quarks, but remains an approximate symmetry useful to consider the deconfinement phase transition ; when quark mass $\rightarrow \infty$ the QCD Lagrangian is invariant under \mathbb{Z}_3 because the mass term \gg kinetic term.

The Polyakov loop: an order parameter for deconfinement

Partition function of QCD in the presence of a static quark Q at position \vec{R} :

$$Z_Q = \int \mathcal{D}A_\mu e^{-S_E(\text{pure gauge})} \times \text{Tr}_c e^{ig \int_0^\beta dx_4 A_4(\vec{R}, x_4)}.$$

One define the Polyakov loop:

$$L(\vec{R}) = \text{Tr}_c e^{ig \int_0^\beta dx_4 A_4(\vec{R}, x_4)} ;$$

it is a color singlet with a \mathbb{Z}_3 charge ($L \rightarrow zL$).

Its thermal expectation value is the so-called Polyakov loop:

$$\Phi(\vec{R}) = \frac{1}{N_c} \langle L(\vec{R}) \rangle_\beta = \frac{Z_Q}{Z_{\text{pureglue}}} = e^{-\beta F_Q(\vec{R})}.$$

F_Q is the free energy associated to a static quark (test charge) added to the pure glue.

Two limit cases:

- $F \rightarrow +\infty$ i.e. there is a confinement of color charges: $\Phi \rightarrow 0$
- $F \rightarrow 0$ i.e. asymptotic freedom: $\Phi \rightarrow 1$

As a consequence, Φ can be seen as an order parameter for the confined ($\Phi \simeq 0$) - deconfined ($\Phi \simeq 1$) phase transition.



It is easy to anticipate that at high temperature, $\Phi \rightarrow 1$ (from the definition of $L(\vec{R})$, $T \rightarrow +\infty$, $\beta \rightarrow 0$ hence the integral and $L(\vec{R}) \simeq 1$).

Pure gauge sector: the effective potential $\mathcal{U}(\Phi, \bar{\Phi}; T)$

\mathbb{Z}_3 linked to confinement is spontaneously broken with temperature. To reproduce this phenomenologically, one can choose a potential (\simeq static gluon pressure term) with this form:

$T < T_0$, Color “confinement”,
 $\langle \Phi \rangle = 0 \longrightarrow$ no \mathbb{Z}_3 breaking

$T > T_0$, Color “deconfinement”,
 $\langle \Phi \rangle \neq 0 \longrightarrow \mathbb{Z}_3$ broken



The PNJL model (Polyakov – Nambu – Jona-Lasinio) in a nutshell

$$\mathcal{L}_{PNJL} = \bar{q} (i\gamma_\mu D^\mu - \hat{m}_0) q + G_1 \left[(\bar{q}q)^2 + (\bar{q}i\gamma_5 \vec{\tau}q)^2 \right] \left(\text{diagram} \simeq \text{diagram} \right) - \mathcal{U}(\Phi[A], \bar{\Phi}[A]; T) \left(\text{diagram} + \text{diagram} \right) + \mu \bar{q} \gamma_0 q$$

* NJL parameters chosen to fit hadronic input in vacuum

* Polyakov loop in imaginary time and Polyakov gauge: $L(\vec{x}) = \mathcal{P} \exp \left[i \int_0^\beta dx_4 A_4(\vec{x}, x_4) \right] \Rightarrow$

Effective field $\Phi = \frac{1}{N_c} \text{Tr}_C L$; L transports the field A_μ from the point in space-time $(\vec{x}, 0)$ to $(\vec{x}, \beta) \Rightarrow \Phi = 0$: confinement; $\Phi = 1$: free propagation (deconfinement)


* Effective potential $\mathcal{U}(\Phi, \bar{\Phi}; T)$ (gluon pressure):

$$\frac{\mathcal{U}(\Phi, \bar{\Phi}; T)}{T^4} = -\frac{b_2(T)}{2} \bar{\Phi} \Phi - \frac{b_3}{6} (\Phi^3 + \bar{\Phi}^3) + \frac{b_4}{4} (\bar{\Phi} \Phi)^2 \text{ and}$$

$$b_2(T) = a_0 + a_1 \left(\frac{T_0}{T} \right) + a_2 \left(\frac{T_0}{T} \right)^2 + a_3 \left(\frac{T_0}{T} \right)^3$$

($T_0 = 270$ MeV)

In the following $\mathcal{U}(\Phi[A], \bar{\Phi}[A]; T)$ can also be chosen in its logarithmic form.



C. Ratti, M. Thaler, W. Weise, hep-ph/0604025 : lattice: O. Kaczmarek, F. Karsch, P. Petreczky, F. Zantow, Phys. Lett. B **543**, 41 (2002).


Mean field grand potential

* **Grand potential at finite temperature and density:** with $E_p = \sqrt{\vec{p}^2 + m^2}$

$$\Omega = \mathcal{U}(\Phi, \bar{\Phi}, T) + \frac{(m - m_0)^2}{2G_1} - 6N_f \int_{\Lambda} \frac{d^3p}{(2\pi)^3} E_p - 2N_f T \int_B \frac{d^3p}{(2\pi)^3} \left\{ \text{Tr}_c \ln \left[1 + \mathbf{L} e^{-(E_p - \mu)/T} \right] + \text{Tr}_c \ln \left[1 + \mathbf{L}^\dagger e^{-(E_p + \mu)/T} \right] \right\}$$

The propagation of the quarks into the medium filled with (background) gluon fields with pressure \mathcal{U} leads to statistical suppression of 1- and 2-quarks propagation (**statistical confinement**):

$$\text{Tr}_c \ln \left[1 + \mathbf{L} e^{-(E_p - \mu)/T} \right] = \ln \left[1 + 3\Phi e^{-\beta(E_p - \mu)} + 3\bar{\Phi} e^{-2\beta(E_p - \mu)} + e^{-3\beta(E_p - \mu)} \right]$$



Cutoff of the model:



The model needs a cutoff in vacuum: \int_0^{Λ} with Λ a typical hadronic scale ($\Lambda \simeq 600$ MeV).

In medium (thermal) part of the model: Boltzmann factors are enough to regularize the integral but ...


pressure with finite and infinite Λ : lack of high momentum quarks to saturate the pressure (Stephan-Boltzmann limit) if $B = \Lambda$.

Effect of the cutoff

* **Consequence on the Polyakov loop:** With $B = +\infty$: “faster” (stronger) crossover ; Also at high density, the deconfinement crossover is larger (“slower” transition) because of the Fermi motion that acts as a large crossover field.





Chiral and deconfinement crossover entanglement



Left: Black: the order parameter $\langle \bar{q}q \rangle$ (chiral symmetry) and Φ (deconfinement) ; Red: deconfinement thermal susceptibility $\chi_\Phi = \frac{\partial \Phi}{\partial T}$; green: susceptibility with m_{fix} (i.e. solving the mean field equations with m fixed to a given value) for seeing the deconfinement transition entanglement.

Right: Φ -susceptibility for different values of μ . The solid lines represent χ_Φ calculated with the Hartree mass, the dashed lines χ_Φ calculated with the constant mass (mean value between the mass at $(T = 0, \mu)$ and m_0). From the green line, we cannot see the dashed lines because χ_Φ is the same if we take a constant mass or the Hartree mass; actually in this region there is no more a chiral transition. The small influence of the mass on Φ allows to disentangle chiral and deconfinement transition.

Deconfinement at high density: mean field phase diagram



⇒ kind of decoupling of deconfinement and chiral restoration at the Critical End Point (CEP).

In this model, 3 phases: chirally broken and “confined” ; chirally broken and “deconfined” ; chirally restored and “deconfined” (QGP)

⇒ tendency to deconfinement at high density (even at vanishing temperature !), an effect driven by the Fermi motion of quarks that broke \mathbb{Z}_3 .

Note that the model is push above its limit (because of the lack of dynamical gluonic degrees of freedom): we take this results as an indication that it *may be* the same in QCD.

Relevance for phenomenology of heavy ion experimental program at high density ?

As a first step, we will take a look on a particular QGP probe, the mesonic correlations.

Link between chiral restoration and deconfinement

Hence one can solve the mean field equation for Φ for any mass (between m_0 and $m_{Hartree}$) without significant changes ; at the contrary, it is wrong for m or the condensate. It depends strongly on Φ : $\Phi \rightarrow 1$ generates more quarks and more screening that breaks the condensate hence a faster chiral restoration.

But it does **not** mean that chiral symmetry restoration cannot occur **before** deconfinement at finite density.

At zero density the Casher argument implies that confinement leads to chiral symmetry breaking. This argument seems valid in the model for a variety of parameters (the quasi-coincidence to a few MeV between chiral restoration and deconfinement is always present).


The Fermi motion plays an important role here: if $\mu > 0$ ($\mu > \mu_{CEP}$) it breaks \mathbb{Z}_3 (it is no more a very good symmetry of the system) \Rightarrow slow increases of Φ that does not influence $\langle \bar{q}q \rangle$ a lot anymore. On the other hand $\mu < \langle q^\dagger q \rangle$ is enough to restore the chiral symmetry.

The two transitions seem to decouple at high density.

Spectral function $Meson \rightarrow q\bar{q}$ as a probe of the phase diagram

Calculation of the mesonic polarisation with the ring approximation (the spectral function describes only $M \rightarrow q\bar{q}$): we choose to remove high momentum quarks ($B = \Lambda$) from the quark loops as a reminder of the QCD asymptotic freedom (high momentum quarks have a weak, negligible interaction for what concern formation of bound states or resonances). Anyway, there is only a small quantitative difference and it is “better” for sum rules (e.g. the V-A sum rule in the vector sector).

No confinement in the PNJL model (only an effect on the thermal bath via the suppression of 1- and 2-quarks Boltzmann factor when $\Phi \simeq 0$): in PNJL “deconfinement” means $\Phi \rightarrow 1$.




Consequence: the (mean field) quark density is vanishing in the vacuum ; saturates SB limit in QGP \Rightarrow deconfinement can be read in quark abundances.

Hence: the $M \rightarrow q\bar{q}$ spectral function will be a tool to study the consequences beyond the mean field of the deconfinement.

Survival of resonant state after the chiral restoration ?


* Qualitative picture

Spectral functions



Confining regime: σ is a narrow resonance ;
chiral symmetry broken
 $\langle \bar{q}q \rangle \neq 0$; $\Phi \simeq 0$


Spectral functions



Deconfinement: σ and π spectral function are
almost the same as an uncorrelated $\bar{q}q$ pair ;
chiral symmetry restored
 $\langle \bar{q}q \rangle \simeq 0$; $\Phi \neq 0$


Survival of resonant state after the chiral restoration ?

Spectral functions



Slow deconfinement (“*Confining*” regime): σ and π still narrow resonances ; chiral symmetry restored
 $\langle \bar{q}q \rangle \simeq 0$; $\Phi \simeq 0$

Evolution on a line at constant μ



The qualitative conclusions


The deconfinement crossover induces very different behavior of the spectral function depending on the density:

- at low density (below the CEP) vanishing of the spectral functions very close to the chiral restoration \Rightarrow “fast” deconfinement that coincides with chiral crossover
- above CEP: vanishing of the spectral function “far” from the chiral transition \Rightarrow “slow” deconfinement that ends well after the 1st order transition.

In the second case: possibility of the existence of a phase where chiral symmetry is restored but the mesons are still confined (no definite answer because of the lack of confinement of the PNJL model but an indication that it may be possible in full QCD).

Quantitative picture



Criterion used to quantify what we can “see” qualitatively: standard deviation of the spectral function. It presents (at moderate density) two plateaux at low temperature (narrow resonance) and high temperature (spectral function $\simeq \text{Im}\Pi_0$).



Left: standard deviation as a function of T for different μ .

Right: σ spectral function for $T = 50$ MeV and two different μ corresponding to the crossing point of the standard deviation: \Rightarrow it is not the absolute value that is used as a criterion to probe the possibility of resonant state, but the relative value with respect to the plateaux.

Comparison with mean field



Left: ending points of the plateaux in the (T, μ) plane.

Right: superposition of the mean field phase diagram and the behavior of the mesons.

\Rightarrow confirms the coherence of the model: the spectral function (the strength) picture follows the order parameter Φ , not the condensate.

As a conclusion

Does this CCS phase really exist (a hadronic spectrum chirally symmetrical composed of true bound state) ? Too soon to know (and impossible to answer within the PNJL model). But at least, this calculation indicates that the phenomenology of the phase transition at high density in future experiment will be very different from the one at zero density:

- obviously because of the 1st order chiral transition
- but also because of this very slow breaking of \mathbb{Z}_3 (indicating a “slow” deconfinement)