

# Dilepton Production with UrQMD

Stephan Endres

Frankfurt Institute for Advanced Studies  
ITP Uni Frankfurt

NeD/TURIC - June 29th, 2012

# Overview

- 1 Introduction
- 2 Dilepton Production in UrQMD
- 3 Results
  - Results for HADES Energies
  - Rho Contribution
  - Cross-sections
- 4 Further Plans
  - Next Steps

# Overview

- 1 Introduction
- 2 Dilepton Production in UrQMD
- 3 Results
  - Results for HADES Energies
  - Rho Contribution
  - Cross-sections
- 4 Further Plans
  - Next Steps

# Overview


- 1 Introduction
- 2 Dilepton Production in UrQMD
- 3 Results
  - Results for HADES Energies
  - Rho Contribution
  - Cross-sections
- 4 Further Plans
  - Next Steps

# Overview

- 1 Introduction
- 2 Dilepton Production in UrQMD
- 3 Results
  - Results for HADES Energies
  - Rho Contribution
  - Cross-sections
- 4 Further Plans
  - Next Steps

# Why Dileptons...?

- Dileptons represent a clean and penetrating probe of hot and dense nuclear matter
- Once produced they do not interact with the surrounding matter
- Aim of studies
  - ⇒ In-medium modification of vector meson properties
  - ⇒ Chiral symmetry restoration



# Dilepton sources in UrQMD

- Dalitz Decays


$\Rightarrow \pi^0, \eta, \eta', \omega, \Delta$

$P \rightarrow \gamma + e^+e^-$

$V \rightarrow P + e^+e^-$

- Direct Decays

$\Rightarrow \rho^0, \omega, \phi$



# Resonances and Branching Ratios in UrQMD

- Two processes possible in UrQMD
  - Collisions**  
(e.g.  $\pi\pi \rightarrow \rho$ )
  - Resonance decays**  
(e.g.  $N^* \rightarrow N + \rho$ )
- At SIS energies, the resonance excitation and decay is dominant
- Branching ratios are in accordance with PDG

Resonance	Mass	Width	$N\pi$	$N\eta$	$N\omega$	$N\rho$
$N_{1440}^*$	1.440	350	0.65			
$N_{1520}^*$	1.515	120	0.60			0.15
$N_{1535}^*$	1.550	140	0.60	0.30		
$N_{1650}^*$	1.645	160	0.60	0.06		0.06
$N_{1675}^*$	1.675	140	0.40			
$N_{1680}^*$	1.680	140	0.60			0.10
$N_{1700}^*$	1.730	150	0.05			0.20
$N_{1710}^*$	1.710	500	0.16	0.15		0.05
$N_{1720}^*$	1.720	550	0.10			0.73
$N_{1900}^*$	1.850	350	0.30	0.14	0.39	0.15
$N_{1990}^*$	1.950	500	0.12			0.43
$N_{2080}^*$	2.000	550	0.42	0.04	0.15	0.12
$N_{2190}^*$	2.150	470	0.29			0.24
$N_{2220}^*$	2.220	550	0.29		0.05	0.22
$N_{2250}^*$	2.250	470	0.18			0.25
$\Delta_{1232}$	1.232	115	1.00			
$\Delta_{1600}^*$	1.700	350	0.10			
$\Delta_{1620}^*$	1.675	160	0.15			0.05
$\Delta_{1700}^*$	1.750	350	0.20			0.25
$\Delta_{1900}^*$	1.840	260	0.25			0.25
$\Delta_{1905}^*$	1.880	350	0.18			0.80
$\Delta_{1910}^*$	1.900	250	0.30			0.10
$\Delta_{1920}^*$	1.920	200	0.27			
$\Delta_{1930}^*$	1.970	350	0.15			0.22
$\Delta_{1950}^*$	1.990	350	0.38			0.08



# Dalitz Decays

- Dalitz decays can be decomposed into the corresponding decays into a virtual photon and the subsequent decay of the photon via electromagnetic conversion

$$\frac{d\Gamma}{dM^2} = \Gamma_{P \rightarrow \gamma \gamma^*, V \rightarrow P \gamma^*} \frac{1}{\pi M^4} M \Gamma_{\gamma^* \rightarrow ee}$$

- Internal conversion probability of the photon

$$M \Gamma_{\gamma^* \rightarrow ee} = \frac{\alpha}{3} M^2 \sqrt{1 - \frac{4m_e^2}{M^2}} \left( 1 + \frac{2m_e^2}{M^2} \right)$$

# Dalitz Decays

- The widths  $\Gamma_{P \rightarrow \gamma \gamma^*}$  and  $\Gamma_{V \rightarrow P \gamma^*}$  can be related to corresponding radiative widths

$$\Gamma_{P \rightarrow \gamma \gamma^*} = 2\Gamma_{P \rightarrow 2\gamma} \left(1 - \frac{M^2}{m_p^2}\right)^3 |F_{P\gamma\gamma^*}(M^2)|^2$$

$$\Gamma_{V \rightarrow P \gamma^*} = 2\Gamma_{V \rightarrow P \gamma} \left[ \left(1 + \frac{M^2}{m_V^2 - m_p^2}\right)^2 - \left(\frac{2m_V M}{m_V^2 - m_p^2}\right)^2 \right]^{3/2} \cdot |F_{VP\gamma^*}(M^2)|^2$$

# Form Factors & Direct Decays

- Form factors for the Dalitz decays are obtained from the vector-meson dominance model (VMD). We use the parametrizations by Landsberg and Li/Ko/Brown/Sorge [L.G. Landsberg, Phys.Rept.128, 301 (1985); C.M. Ko et al., Nucl.Phys. A610, 342C (1996)]
- The width for the direct decay of a vector meson  $V$  to a dilepton pair varies with the dilepton mass like  $M^{-3}$

$$\Gamma_{V \rightarrow ee}(M) = \frac{\Gamma_{V \rightarrow ee}(m_V)}{m_V} \frac{m_V^4}{M^3} \sqrt{1 - \frac{4m_e^2}{M^2}} \left(1 + \frac{2m_e^2}{M^2}\right)$$


# Time Integration Method (Shining)

- Shining in UrQMD applied for  $\Delta$ ,  $\rho$ ,  $\omega$ ,  $\phi$  and  $\eta'$
- *Assumption*: Resonance can continuously emit dileptons over its whole lifetime
- Integration of the dilepton emission rate over time
- Collisional broadening of each individual parent resonance is taken into account

$$\frac{dN_{ee}}{dM} = \frac{\Delta N_{ee}}{\Delta M} = \sum_{j=1}^{N_{\Delta M}} \int_{t_i^j}^{t_f^j} \frac{dt}{\gamma} \frac{\Gamma_{ee}(M)}{\Delta M}$$


# p + p @ 1.25 GeV

- $E_{lab} = 1.25$  GeV just below  $\eta$  threshold
- Small sub-threshold contribution from  $\rho$  expected
- Good agreement with data at low masses
- Too many dileptons from 0.3 GeV on  $\rightarrow$  especially  $\rho^0$





# p + p @ 2.2 GeV

- Above  $\eta$  threshold, energy sufficient to reach pole mass of  $\omega$  and  $\rho$
- Overestimation of  $\rho$  contribution even larger than at 1.25 GeV




# $p_t$ Spectra for $p + p$ @ 2.2 GeV



- $p_t$  spectra are described well, especially in the low mass region dominated by the  $\pi^0$  Dalitz decay
- We again see an overestimation of the  $\rho$  meson contribution

# $p_t$ Spectra for $p + p$ @ 2.2 GeV

- In the intermediate mass region we see a our Delta contribution overshooting the data in for high  $p_t$







# n + p @ 1.25 GeV

- Deuteron beam with 1.25A GeV has been used by HADES besides p+p
  - Trigger on forward-going protons in order to select the (quasi-free) np collisions
  - Fermi motion of the bound nucleons in the deuteron leads to a smearing of the NN collision energy → reaching above  $\eta$ -production threshold
- ⇒ One can not easily compare data with pure n+p simulations

# $n + p @ 1.25 \text{ GeV}$




- $\eta$  and more  $\rho$  are produced in  $d+p$ , compared to  $n+p$
- However, even for  $d+p$  the Yield is underestimated by a factor of about 5 to 10

# Study of A+A collisions



- In nucleus-nucleus collisions, additional effects compared to pp are expected
  - Fermi Momentum
  - Not only p+p, but also p+n and n+n collisions
  - Secondary interactions, depending on system size and energy
- Vector meson spectral functions may be changed in the medium
  - Shift of the pole mass (of the  $\rho$ )
  - Resonance melting in the medium
- In UrQMD, no such in-medium effects are implemented

# C + C @ 1 AGeV

- Here the  $\rho$  contribution fits quite well
- Underestimation below for energies above the pion peak
- Hardly any  $\omega$  produced,  $\eta$  is relatively small




# Ar + KCl @ 1.76 AGeV & C + C @ 2 AGeV




- As in elementary reactions, we get too many dileptons via the  $\rho^0$  resonance, especially in the high-mass tail
- How are the  $\rho$  mesons produced?

# $\rho^0$ Contribution



- Main contribution below pole mass by  $N^*_{1520}$  resonance
- For pole mass peak via  $N^*_{1680}$ ,  $N^*_{1680}$  and  $N^*_{1700}$
- Other sources negligible

# $\rho^0$ Contribution




- Main contribution below pole mass again by  $N^*_{1520}$
- For pole mass peak via  $N^*_{1720}$ ,  $\Delta^*_{1700}$  and  $\Delta^*_{1905}$
- $\pi\pi \rightarrow \rho$  makes up 10%

## Why too many dileptons from $\rho$ ?

- All HADES energies are close to thresholds
  - Cross-sections change rapidly with small energy differences
  - Are the cross-sections in order?
- Do in-medium effects not included in UrQMD play a role ?
- Possible  $\sigma$  channel for a part of what we treat as a  $\rho$ ?
- Why do we see the large overestimation for invariant masses above 700 MeV (pole mass)?




# Cross-sections $N+N \rightarrow \rho^0 + X$



- Good description of inclusive cross-section
- For energies near threshold,  $\sigma$  might be too high, but no data are available for this region
- However, can't explain the overestimation at masses higher than the  $\rho$  pole mass

# Cross-sections $\pi+N \rightarrow \rho+X$

- $\rho^+$  production overestimated below  $\sqrt{s} = 3$  GeV (but not relevant for dileptons)
- $\rho^0$  from UrQMD fits quite well to data, except for threshold region
- Does experiment just don't see these  $\rho$  which are significantly below pol mass?



# Cross-sections

- Other cross-sections might as well play a role, especially R+N or R+R
  - Not relevant for elementary, but significant for A+A collision
  - Little data is available for these cases
- ⇒ Has to be checked next...

- Reduce the overestimation dilepton production in UrQMD, especially via the  $\rho^0$ ,  $\Delta$  and  $\eta$  channel
- Coarse graining to be done for HADES energies
  - \* Take local temperature and baryon chemical potential as functions of space and time
  - \* Accumulate an ensemble of events and determine local variables via coarse graining
- Dilepton calculation with hybrid model (transport + hydro)
  - \* Previous work (Dimuons from NA60) by Elvira Santini

[E. Santini et al., Phys.Rev. C84, 014901 (2011)]
- \* Proceed with this work and calculate yields for RHIC and LHC energies