On the **Gauge Boson's Properties** in a **Two-Color Theory** with **Two Adjoint Flavors**

Axel Maas



February 2012 Strong Interactions beyond the Standard Model Bad Honnef Germany

On the Gauge Boson's Properties in Minimal Walking Technicolor

Axel Maas



February 2012 Strong Interactions beyond the Standard Model Bad Honnef Germany

Set-up and problems

- Set-up and problems
- Describing elementary particles

- Set-up and problems
- Describing elementary particles
 - Correlation functions

[Maas 1106.3942]

- Set-up and problems
- Describing elementary particles
 - Correlation functions

[Maas 1106.3942]

• Gauge bosons in minimal walking technicolor

- Set-up and problems
- Describing elementary particles
 - Correlation functions

[Maas 1106.3942]

- Gauge bosons in minimal walking technicolor
 - Propagators
 - Vertices
 - Zero mass limit

[Maas JHEP '11]

• Summary

Describe the standard model and beyond

- Describe the standard model and beyond
- 12+ orders of magnitude of scales
 - Neutrino mass to new physics
 - Can be intricately related (e.g. see-saw)

- Describe the standard model and beyond
- 12+ orders of magnitude of scales
 - Neutrino mass to new physics
 - Can be intricately related (e.g. see-saw)
- Chiral theory

- Describe the standard model and beyond
- 12+ orders of magnitude of scales
 - Neutrino mass to new physics
 - Can be intricately related (e.g. see-saw)
- Chiral theory
- Strong interactions
 - QCD, technicolor,...

- Describe the standard model and beyond
- 12+ orders of magnitude of scales
 - Neutrino mass to new physics
 - Can be intricately related (e.g. see-saw)
- Chiral theory
- Strong interactions
 - QCD, technicolor,...
 - In general non-factorizable

[Andersen et al. EPJ+ 2011]

• Here: Technicolor extension of the standard model

- Here: Technicolor extension of the standard model
 - "Minimal walking technicolor"
 - SU(2), 2 flavors of massless adjoint fermions

- Here: Technicolor extension of the standard model
 - "Minimal walking technicolor"
 - SU(2), 2 flavors of massless adjoint fermions
 - Likely walking
 - Slow evolution with momenta
 - Many orders of magnitude
 - Large scale compared to the standard model

- Here: Technicolor extension of the standard model
 - "Minimal walking technicolor"
 - SU(2), 2 flavors of massless adjoint fermions
 - Likely walking
 - Slow evolution with momenta
 - Many orders of magnitude
 - Large scale compared to the standard model
 - Light, chiral fermions

- Here: Technicolor extension of the standard model
 - "Minimal walking technicolor"
 - SU(2), 2 flavors of massless adjoint fermions
 - Likely walking
 - Slow evolution with momenta
 - Many orders of magnitude
 - Large scale compared to the standard model
 - Light, chiral fermions
 - Strongly interacting chiral gauge theory when coupled to the standard model

No gauge-invariant elementary states

- No gauge-invariant elementary states
 - Includes elementary or composite Higgs
 - Nielsen identities only imply gauge-parameter invariance of mass, but not gauge invariance

- No gauge-invariant elementary states
 - Includes elementary or composite Higgs
 - Nielsen identities only imply gauge-parameter invariance of mass, but not gauge invariance
 - Electron requires photon cloud

- No gauge-invariant elementary states
 - Includes elementary or composite Higgs
 - Nielsen identities only imply gauge-parameter invariance of mass, but not gauge invariance
 - Electron requires photon cloud
- Only gauge-invariant quantities are bound states and cross sections of bound states

- No gauge-invariant elementary states
 - Includes elementary or composite Higgs
 - Nielsen identities only imply gauge-parameter invariance of mass, but not gauge invariance
 - Electron requires photon cloud
- Only gauge-invariant quantities are bound states and cross sections of bound states
- Gauge-invariant bound states of tops, Ws, Higgses... may be rather unstable

 Description in terms of elementary degrees of freedom rather successful

- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory

- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory
 - PDG data for Higgs etc. based on the elementary, gauge-dependent degrees of freedom

- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory
 - PDG data for Higgs etc. based on the elementary, gauge-dependent degrees of freedom
 - Gauge-dependent intermediate step

- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory
 - PDG data for Higgs etc. based on the elementary, gauge-dependent degrees of freedom
 - Gauge-dependent intermediate step
 - Correlation functions contain all information
 - Masses, running couplings, widths,...

- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory
 - PDG data for Higgs etc. based on the elementary, gauge-dependent degrees of freedom
 - Gauge-dependent intermediate step
 - Correlation functions contain all information
 - Masses, running couplings, widths,...
 - Final quantities are gauge-invariant


- Description in terms of elementary degrees of freedom rather successful
 - All of perturbation theory
 - PDG data for Higgs etc. based on the elementary, gauge-dependent degrees of freedom
 - Gauge-dependent intermediate step
 - Correlation functions contain all information
 - Masses, running couplings, widths,...
 - Final quantities are gauge-invariant
- Non-perturbative extension
 - Correlation functions central element

[Maas 1106.3942 Fischer JPG 2006]


[Maas 1106.3942 Fischer JPG 2006]



[Maas 1106.3942 Fischer JPG 2006]




[Maas 1106.3942 Fischer JPG 2006]




[Maas 1106.3942 Fischer JPG 2006]


• A theory can be described in terms of correlation functions




Quantization




[Maas 1106.3942 Fischer JPG 2006]




[Maas 1106.3942 Fischer JPG 2006]




[Maas 1106.3942 Fischer JPG 2006]




[Maas 1106.3942 Fischer JPG 2006]



[Maas 1106.3942 Fischer JPG 2006]



[Maas 1106.3942 Fischer JPG 2006]



Physics			

Physics	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control

•					
Physics	All elementary	Chiral, light fermions	Strong interactions	Different scales	Control
Methods	particles				

1

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good
Funct. Equat.	Good	Good	Good	Good	Difficult

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good
Funct. Equat.	Good	Good	Good	Good	Difficult
Models	Impossible	Good	Good	Difficult	Difficult

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good
Funct. Equat.	Good	Good	Good	Good	Difficult
Models	Impossible	Good	Good	Difficult	Difficult

Combination of multiple methods enhances advantages


Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good
Funct. Equat.	Good	Good	Good	Good	Difficult
Models	Impossible	Good	Good	Difficult	Difficult

- Combination of multiple methods enhances advantages
- Lattice and functional method combination well tested in QCD

Physics Methods	All elementary particles	Chiral, light fermions	Strong interactions	Different scales	Control
PT	Good	Good	Impossible	Good	Good
Lattice	Good	Difficult	Good	Difficult	Good
Funct. Equat.	Good	Good	Good	Good	Difficult
Models	Impossible	Good	Good	Difficult	Difficult

- Combination of multiple methods enhances advantages
- Lattice and functional method combination well tested in QCD
- Perturbation theory covered in lattice and functional equations


• Lattice



Lattice


- Discretize space-time in a box and calculate the pathintegral and expectation values explicitly
- ✓Full non-perturbative dynamics correctly implemented
- Finite volume artifacts, disparate scales most severe obstacles

- Lattice
 - Discretize space-time in a box and calculate the pathintegral and expectation values explicitly
 - Full non-perturbative dynamics correctly implemented
 - Finite volume artifacts, disparate scales most severe obstacles
- Functional methods (DSE, RGE...)




- Lattice
 - Discretize space-time in a box and calculate the pathintegral and expectation values explicitly
 - Full non-perturbative dynamics correctly implemented
 - Finite volume artifacts, disparate scales most severe obstacles
- Functional methods (DSE, RGE...)
 - Coupled non-linear integral equations must be solved
 Continuum, infinite volume, analytically accessible
 Kequires (often uncontrolled) approximations

- Lattice
 - Discretize space-time in a box and calculate the pathintegral and expectation values explicitly
 - Full non-perturbative dynamics correctly implemented
 - Finite volume artifacts, disparate scales most severe obstacles
- Functional methods (DSE, RGE...)
 - Coupled non-linear integral equations must be solved
 Continuum, infinite volume, analytically accessible
 Kequires (often uncontrolled) approximations
- Perturbation theory included




[Maas 1106.3942 Fischer JPG 2006]

• Standard approach: One method – One result




[Maas 1106.3942 Fischer JPG 2006]

• Standard approach: One method – One result



[Maas 1106.3942 Fischer JPG 2006]

• Standard approach: One method – One result




Method 2




[Maas 1106.3942 Fischer JPG 2006]

Lattice calculations





Functional calculations


[Maas 1106.3942 Fischer JPG 2006]




Functional calculations











[Maas 1106.3942 Fischer JPG 2006]



Lattice calculations

Functional calculations

[Maas 1106.3942 Fischer JPG 2006]



Lattice calculations

• Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]

- Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]
- Bound-state properties [Fischer JPG 2006]

- Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]
- Bound-state properties [Fischer JPG 2006]
- Finite temperature and density
 - Phase diagram of Yang-Mills theory [Pawlowski et al., Maas et al.,...]
 - Phase diagram of (quenched) QCD

[Pawlowski et al., Fischer et al., Nickel et al.,...]

- Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]
- Bound-state properties [Fischer JPG 2006]
- Finite temperature and density
 - Phase diagram of Yang-Mills theory [Pawlowski et al., Maas et al.,...]
 - Phase diagram of (quenched) QCD
- Higgs physics [Maas EPJC 10]

Applications of this framework

- Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]
- Bound-state properties [Fischer JPG 2006]
- Finite temperature and density
 - Phase diagram of Yang-Mills theory [Pawlowski et al., Maas et al.,...]
 - Phase diagram of (quenched) QCD [Pawlowski et al., Fischer et al., Nickel et al.,...]
- Higgs physics [Maas EPJC 10]
- Many works based on model gauge bosons

[Roberts, Tandy et al. 2012, Alkofer & von Smekal PR 2001]

Applications of this framework

- Yang-Mills theory and QCD [Maas 1106.3942, Fischer JPG 2006]
- Bound-state properties [Fischer JPG 2006]
- Finite temperature and density
 - Phase diagram of Yang-Mills theory [Pawlowski et al., Maas et al.,...]
 - Phase diagram of (quenched) QCD [Pawlowski et al., Fischer et al., Nickel et al.,...]
- Higgs physics [Maas EPJC 10]
- Many works based on model gauge bosons [Roberts, Tandy et al. 2012, Alkofer & von Smekal PR 2001]
- Here: Minimal walking technicolor

[Sannino 2009, Maas JHEP 2011]

• Configurations by Del Debbio, Lucini, Patella, Pica, Rago

[Maas 1106.3942 Fischer JPG 2006]



- Pure technicolor sector
 - SU(2)+2 adjoint flavors of techniquarks


- Pure technicolor sector
 - SU(2)+2 adjoint flavors of techniquarks
- Massless limit relevant but what is the mass?
 - Scale setting without experimental input complicated

- Pure technicolor sector
 - SU(2)+2 adjoint flavors of techniquarks
- Massless limit relevant but what is the mass?
 - Scale setting without experimental input complicated
 - No scale if the theory is conformal
 - Well-defined if the theory is quasi-conformal


- Pure technicolor sector
 - SU(2)+2 adjoint flavors of techniquarks
- Massless limit relevant but what is the mass?
 - Scale setting without experimental input complicated
 - No scale if the theory is conformal
 - Well-defined if the theory is quasi-conformal
- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass

- Pure technicolor sector
 - SU(2)+2 adjoint flavors of techniquarks
- Massless limit relevant but what is the mass?
 - Scale setting without experimental input complicated
 - No scale if the theory is conformal
 - Well-defined if the theory is quasi-conformal
- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass
- Other possibilities exist
 - Lattice scale, mesons, fixed scale,...


- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass



- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass




- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass
 - Compatible with string tension



- Scale setting by a scale with smooth limit to the Yang-Mills case
 - Techniglueball mass
 - Compatible with string tension
 - Scale quickly growing towards massless fermions


[Maas 1106.3942 Fischer JPG 2006]

Quantization




Quantization and gauge fixing


• Gauge freedom: Choice of internal coordinate system




- Gauge freedom: Choice of internal coordinate system
- Particle fields change under gauge transformations
 - So do most correlation functions



- Gauge freedom: Choice of internal coordinate system
- Particle fields change under gauge transformations
 - So do most correlation functions
- Requires a choice of gauge/coordinate system




- Gauge freedom: Choice of internal coordinate system
- Particle fields change under gauge transformations
 - So do most correlation functions
- Requires a choice of gauge/coordinate system
- Indirect by a condition on the gauge fields
- Local condition in perturbation theory



- Gauge freedom: Choice of internal coordinate system
- Particle fields change under gauge transformations
 - So do most correlation functions
- Requires a choice of gauge/coordinate system
- Indirect by a condition on the gauge fields
- Local condition in perturbation theory
 - Used here: Landau gauge $\partial^{\mu}A^{a}_{\mu}=0$


Quantization and gauge fixing

[Maas 1106.3942]




Quantization and gauge fixing

• Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]




- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$




- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$


- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$
 - Gribov-Singer ambiguity



- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$
 - Gribov-Singer ambiguity: Requires non-local constraints




- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$
 - Gribov-Singer ambiguity: Requires non-local constraints
- Resolution by specifying a sampling procedure for Gribov copies [Maas PLB 10/PRD 09, Fischer, Maas, Pawlowski AoP 09, Neuberger PLB '87, von Smekal et al. '08/'09]




- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$
 - Gribov-Singer ambiguity: Requires non-local constraints
- Resolution by specifying a sampling procedure for Gribov copies [Maas PLB 10/PRD 09, Fischer, Maas, Pawlowski AoP 09, Neuberger PLB '87, von Smekal et al. '08/'09]
- Well-defined gauge prescription
 - Necessary for comparison of methods
 - Gives a precise meaning for the correlation functions
 - Here: Minimal (or average) Landau gauge

- Insufficient beyond perturbation theory [Gribov NPA 77, Singer CMP 78]
 - Gribov copies: More than one solution to $\partial^{\mu} A^{a}_{\mu} = 0$
 - Gribov-Singer ambiguity: Requires non-local constraints
- Resolution by specifying a sampling procedure for Gribov copies [Maas PLB 10/PRD 09, Fischer, Maas, Pawlowski AoP 09, Neuberger PLB '87, von Smekal et al. '08/'09]
- Well-defined gauge prescription
 - Necessary for comparison of methods
 - Gives a precise meaning for the correlation functions
 - Here: Minimal (or average) Landau gauge
- Impact of Gribov-Singer ambiguity very similar to Yang-Mills theory at same lattice size/glueball mass [Maas JHEP 2011]
 - Similar to other confining theories [Maas EPJC 2010]


[Maas 1106.3942 Fischer JPG 2006]



Quantization




[Maas 1106.3942 Fischer JPG 2006]




Propagators $\langle A^a_{\mu} A^b_{\nu} \rangle$

Quantization

 $D_{\mu\nu}^{ab}(x-y) = \langle A_{\mu}^{a}(x) A_{\nu}^{b}(y) \rangle$





[Maas 1106.3942 Fischer JPG 2006]




Propagators $\langle A^a_{\mu} A^b_{\nu} \rangle$








$$D_{\mu\nu}^{ab}(p) = \delta^{ab}(\delta_{\mu\nu} - \frac{p_{\mu}p_{\nu}}{p^{2}})D(p)$$




Rescaled Yang-Mills results for comparison

Technigluon propagator




- Rescaled Yang-Mills results for comparison
- Very similar to Yang-Mills result

Technigluon propagator



- Rescaled Yang-Mills results for comparison
- Very similar to Yang-Mills result

Technigluon propagator




- Rescaled Yang-Mills results for comparison
- Very similar to Yang-Mills result

[Maas JHEP 2011]

Technigluon propagator


Technigluon propagator



Rescaled Yang-Mills results for comparison

 Very similar to Yang-Mills result even close to massless limit


Technigluon propagator



- Rescaled Yang-Mills results for comparison
- Very similar to Yang-Mills result even close to massless limit
 - Similar to QCD, scalar theories [Maas 1106.3942]

Technigluon propagator


Technigluon propagator





- Rescaled Yang-Mills results for comparison
- Very similar to Yang-Mills result even close to massless limit
 - Similar to QCD, scalar theories [Maas 1106.3942]
- Faddeev-Popov ghost propagator similar


Describing a theory


[Maas 1106.3942 Fischer JPG 2006]












[Maas JHEP 2011]



Significant structure for Yang-Mills theory

[Maas JHEP 2011]

Three-technigluon vertex 2 B(q,0,q) $am_0=0.5 at 16 \times 8^3 a^{-1}=1.7 TeV$ Yang-Mills at 64 imes 24 3 a⁻¹=1.5 TeV 0 Yang-Mills at 64 imes 24³ a⁻¹=7.2 TeV 1.5 Ę ф 0.5 -0.5 15 10 20 25 5 q [TeV]

• Significant structure for Yang-Mills theory

• Similar at finite mass

[Maas JHEP 2011]


Three-technigluon vertex 2 B(q,0,q) am,=0.5 at 16×8^3 a⁻¹=1.7 TeV am_=-1.00 at 24×12^3 a⁻¹=3.0 TeV 1.5 Yang-Mills at 64 × 24³ a⁻¹=1.5 TeV 0 Yang-Mills at 64 × 24³ a⁻¹=7.2 TeV 0.5 0 -0.5 15 20 10 25 5 q [TeV]

Significant structure for Yang-Mills theory

• Similar at finite mass

[Maas JHEP 2011]


Three-technigluon vertex



- Significant structure for Yang-Mills theory
- Similar at finite mass

[Maas JHEP 2011]


Three-technigluon vertex



- Significant structure for Yang-Mills theory
- Similar at finite mass
- Flattens out towards chiral limit?

[Maas JHEP 2011]

Three-technigluon vertex




- Significant structure for Yang-Mills theory
- Similar at finite mass
- Flattens out towards chiral limit?
- Strongly affected by lattice asymmetry


[Maas JHEP 2011]

• Running coupling computed from correlation functions [von Smekal et al. AoP 1997]


- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]




- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]
- Turnover in Yang-Mills case depends on the scheme [Pawlowski, Maas, Fischer AoP 2009, Maas 1106.3942]




- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]
- Turnover in Yang-Mills case depends on the scheme [Pawlowski, Maas, Fischer AoP 2009, Maas 1106.3942]



- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]
- Turnover in Yang-Mills case depends on the scheme [Pawlowski, Maas, Fischer AoP 2009, Maas 1106.3942]



- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]
- Turnover in Yang-Mills case depends on the scheme [Pawlowski, Maas, Fischer AoP 2009, Maas 1106.3942]



- Running coupling computed from correlation functions [von Smekal et al. AoP 1997]
 - MiniMOM scheme, related to Msbar [von Smekal et al. PLB 2008]
- Turnover in Yang-Mills case depends on the scheme [Pawlowski, Maas, Fischer AoP 2009, Maas 1106.3942]
- Flattens out towards chiral limit

 Elementary degrees of freedom gaugedependent

- Elementary degrees of freedom gaugedependent
- Gauge-fixed correlation functions useful
 - Intermediate steps to gauge-invariant physics
 - Combination of methods

- Elementary degrees of freedom gaugedependent
- Gauge-fixed correlation functions useful
 - Intermediate steps to gauge-invariant physics
 - Combination of methods
- Gauge bosons in minimal walking technicolor
 - Outside massless limit similar to Yang-Mills case
 - Massless limit subtle

- Elementary degrees of freedom gaugedependent
- Gauge-fixed correlation functions useful
 - Intermediate steps to gauge-invariant physics
 - Combination of methods
- Gauge bosons in minimal walking technicolor
 - Outside massless limit similar to Yang-Mills case
 - Massless limit subtle
- Next: Fermion properties