

Phase diagrams by strong coupling methods: QCD at finite temperature and density


Owe Philipsen



- Introduction: The QCD phase diagram
- The deconfinement transition in Yang-Mills theory
- The deconfinement transition in QCD with heavy quarks

in collaboration with M. Fromm, J. Langelage, S. Lottini

The (lattice) calculable region of the phase diagram




- Sign problem prohibits direct simulation, circumvented by approximate methods: **reweighting, Taylor expansion, imaginary chem. pot., need** $\mu/T \lesssim 1$ ($\mu = \mu_B/3$)
- Upper region: equation of state, screening masses, quark number susceptibilities etc. under control
- Here: phase diagram itself, so far based on models, **most difficult!**

The good news: comparing $T_c(\mu)$

de Forcrand, Kratochvila 05


$N_t = 4, N_f = 4$; same actions (unimproved staggered), same mass



imaginary μ
 2 param. imag. μ
 dble reweighting, LY zeros
 Same, susceptibilities
 canonical

Agreement for $\mu/T \lesssim 1$

The order of the p.t., arbitrary quark masses $\mu = 0$



deconfinement p.t.:
breaking of global $Z(3)$ symmetry

deconfinement critical line

chiral critical line


chiral p.t.

restoration of global symmetry in flavour space

$$SU(2)_L \times SU(2)_R \times U(1)_A$$

↑
anomalous

Much harder: is there a QCD critical point?




Two strategies:

1 follow **vertical line**: $m = m_{\text{phys}}$, turn on μ

2 follow **critical surface**: $m = m_{\text{crit}}(\mu)$

Some methods trying (1) give indications of critical point, but systematics not yet controlled

➔ On coarse lattice exotic scenario:
no chiral critical point at small density



Weakening of p.t. with chemical potential also for:

-Heavy quarks

-Light quarks with finite isospin density

-Electroweak phase transition with finite lepton density

Fromm, Langelage, Lottini, O.P. 11

Kogut, Sinclair 07

Gynther 03

Larger densities? Try effective theories!

- Example e.w. phase transition: success with dimensional reduction!
- Scale “separation”
Integrate hard scale perturbatively, treat eff. 3d theory on lattice,
valid for sufficiently weak coupling $g^2 T < gT < 2\pi T$
- Does **not** work for the QCD transition, breaks $Z(3)$ symmetry of Yang-Mills theory
- Bottom up construction of $Z(N)$ -invariant theory by matching couplings:
works for $SU(2)$, not for $SU(3)$
Vuorinen, Yaffe; de Forcrand, Kurkela;
- Here: solution by strong coupling expansion!
- Convergent series within finite convergence radius, valid in confined phase

Starting point: Wilson's lattice action

Partition function; link variables as degrees of freedom

$$Z = \int \prod_{x,\mu} dU(x; \mu) \exp(-S_{YM}) \equiv \int DU \exp(-S_{YM})$$

Wilson's gauge action

$$S_W = -\frac{\beta}{N} \sum_p \text{ReTr}(U_p) = \sum_p S_p \quad \beta = \frac{2N}{g^2}$$

Plaquette:

$$\square \rightarrow 1 + ia^2 g F_{\mu\nu} - \frac{a^4 g^2}{2} F_{\mu\nu} F^{\mu\nu} + O(a^6) + \dots$$

$U_\mu(x) = e^{-ia g A_\mu(x)}$

$$T = \frac{1}{aN_t} \quad \text{continuum limit} \quad a \rightarrow 0, N_t \rightarrow \infty$$

Small $\beta(a) \Rightarrow$ small T

The strong coupling expansion

Expansion in irreducible characters $\chi_r(U) = \text{Tr} D_r(U)$

$$\exp(-S_p) = c_0(\beta) \left\{ 1 + \sum_{r \neq 0} d_r c_r(\beta) \chi_r(U_p) \right\}$$

Expansion parameters $c_r(\beta)$ are combinations of modified Bessel functions (for $SU(N)$)

$$c_f \equiv u \sim \beta + \dots$$

$$c_{ad} \sim \beta^2 + \dots$$

Higher dimensional representations go with higher orders in β

Here: effective lattice theory, general strategy

- Start with the partition function of (3+1) dimensional lattice gauge field theory at finite temperature
- Integrate out degrees of freedom in order to have an effective action in terms of the order parameter (*here*: Polyakov loop)

$$-S_{eff} = \lambda_1 S_1 + \lambda_2 S_2 + \lambda_3 S_3 + \dots$$

- S_n depend only on Polyakov loops
 - Find critical parameters $\lambda_{n,crit}$ and relate back to critical lattice couplings β_{crit} for different N_τ
- Crucial to know mappings $\lambda_n(N_\tau, \beta)$

The effective theory for SU(2)

- Split temporal and spatial link integration and use character expansion ($c_r(\beta)$: expansion parameter of representation r)

$$Z = \int [dW] \exp \left\{ \ln \int [dU_i] \prod_p \left[1 + \sum_{r \neq 0} d_r c_r(\beta) \chi_r(U_p) \right] \right\}$$
$$\equiv \int [dW] \exp [-S_{eff}] \quad W(\vec{X}) = \prod_{\tau=1}^{N_\tau} U_0(\tau, \vec{X})$$


Integration rule 1

$$\int dU \chi_r(U) = \delta_{r,0}$$

Integration rule 2

$$\int dU \chi_r(UV) \chi_s(U^{-1}W) = \delta_{rs} \frac{1}{d_r} \chi_r(VW)$$

Used to perform the occurring group integrations



- Leading order graph in case of $N_\tau = 4$:



Figure: 4 plaquettes in fundamental representation lead to a 2 Polyakov loop interaction term

- Integration of spatial link variables leads to

$$-S_1 = u^{N_\tau} \sum_{\langle ij \rangle} \text{tr } W_i \text{tr } W_j$$

- Possible generalizations: larger distance, higher dimensional representations, larger number of loops involved, ...
- *Here*: Decorate LO graph with additional spatial and temporal plaquettes

- $Z(2)$ symmetric 3 dimensional partition function

$$Z = \int [dW] \exp \left[\lambda_1 \sum_{\langle ij \rangle} \text{tr } W_i \text{tr } W_j \right]$$

- Can be further simplified by using $L \equiv \text{tr } W$ as degrees of freedom: ordinary integration instead of group integration
- Introduces potential term: $V_{SU(2)} = \frac{1}{2} \sum_i \ln [4 - L_i^2]$

$$Z = \int [dL] \exp \left[\lambda_1 \sum_{\langle ij \rangle} L_i L_j + \frac{1}{2} \sum_i \ln [4 - L_i^2] \right]$$

$$\lambda_1(u, N_\tau \geq 5) = u^{N_\tau} \exp \left[N_\tau \left(4u^4 - 4u^6 + \frac{140}{3}u^8 - \frac{36044}{405}u^{10} \right) \right]$$

- **One** determination of $\lambda_{1,crit}$ gives all $\beta_{crit}(N_\tau)$

- Subclass of higher order interaction terms (Powers of the leading order term) arrange schematically as

$$-S_{eff} = \lambda_1(LL) - \frac{\lambda_1^2}{2}(LL)^2 + \frac{\lambda_1^3}{3}(LL)^3 - \dots = \ln \left[1 + \lambda_1(LL) \right]$$


- SU(2) effective theory to be simulated

$$Z = \int [dL] \prod_i \sqrt{4 - L_i^2} \prod_{\langle ij \rangle} \left[1 + \lambda_1 L_i L_j \right]$$

Generalisation to SU(3)

- SU(3) straightforward, but: Now also with anti-fundamental representation (i.e. L_i are complex)

$$\begin{aligned} Z &= \int [dL] \exp [-S_1 + V_{SU(3)}] \\ &= \int [dL] \prod_{\langle ij \rangle} \left[1 + 2\lambda_1 \operatorname{Re}(L_i L_j^*) \right] * \\ &\quad * \prod_i \sqrt{27 - 18|L_i|^2 + 8\operatorname{Re}L_i^3 - |L_i|^4} \end{aligned}$$



- Functional form of $\lambda_1(N_\tau, u)$ and next-to-nearest-neighbour effects are analogous to SU(2)

Numerical evaluation of effective theories

Monte Carlo simulation of scalar model, Metropolis update

Search for criticality:

Binder cumulant: $B(|L|) = 1 - \frac{\langle |L|^4 \rangle}{3\langle |L|^2 \rangle^2} \rightarrow \lambda_{1,c}(N_s)$ is the minimum


Susceptibility: $\chi(|L|) = \langle (|L| - \langle |L| \rangle)^2 \rangle \rightarrow \lambda_{1,c}(N_s)$ is the maximum

Finite size scaling: $\lambda_{1,c}(N_s) = \lambda_{1,c} + bN_s^{-1/\nu}$



$\nu = 1/3$ for the 1st order $SU(3)$, ν_{Ising3D} for $SU(2)$.



Typical sizes range from $N_s = 6$ to 16; time needed is of order a **few days** on an ordinary PC.

Numerical results for SU(3)



Order-disorder transition







First order phase transition for SU(3) in the thermodynamic limit!



The influence of a second coupling

NLO-couplings: next-to-nearest neighbour, adjoint rep. loops



...gets **very** small for large N_τ !

Numerical results for SU(2), one coupling




Second order (3d Ising) phase transition for SU(2) in the thermodynamic limit!

Mapping back to 4d finite T Yang-Mills

Inverting

$$\lambda_1(N_\tau, \beta) \rightarrow \beta_c(\lambda_{1,c}, N_\tau) \quad \dots \text{points at reasonable convergence}$$




SU(3)


Comparison with 4d Monte Carlo

Relative accuracy for β_c compared to the full theory

SU(2)




SU(3)



Note: influence of some couplings checked explicitly!

Extrapolation to continuum limit!



-error bars: difference between last two orders in strong coupling exp.

-using non-perturbative beta-function (4d T=0 lattice)

-all data points from one single 3d MC simulation!

Including fermions

N_f (degenerate) fermions $\implies S = S_{\text{gauge}} + S_q[U, \psi, \bar{\psi}]$

$$S_q = \sum_{x,y;f} \bar{\psi}_{f,y} \left(\mathbb{1} - \kappa H[U] \right)_{yx} \psi_{f,x} \quad , \quad H[U]_{yx} = \sum_{\pm\mu} \delta_{y,x+\hat{\mu}} (\mathbb{1} + \gamma_{\mu}) U_{x,\mu}$$


Integrate the Grassmann variables $\psi, \bar{\psi}$:

$$S = S_{\text{gauge}} - N_f \text{Tr} \log(\mathbb{1} - \kappa H)$$

Expand in the *hopping parameter* $\kappa = 1/(2aM + 8)$: [*]

$$S = S_{\text{gauge}} + N_f \sum_{\ell=1}^{\infty} \frac{\kappa^{\ell}}{\ell} \text{Tr} H[U]^{\ell}$$


Links along imaginary time gain $\exp(\pm\mu a)$



reabsorbed in gauge part: $\begin{cases} \beta \rightarrow \beta + \mathcal{O}(\kappa^4) \\ u(\beta) \rightarrow u(\beta, \kappa) \end{cases}$



LO Polyakov "magnetic" term $\sim \begin{cases} \underbrace{(2\kappa e^{+a\mu})^{N_\tau}}_{h_1} L \\ \underbrace{(2\kappa e^{-a\mu})^{N_\tau}}_{\bar{h}_1} L^* \end{cases}$



higher corrections to the above:

$$h_1 = (2\kappa e^{a\mu})^{N_\tau} \left[1 + \mathcal{O}(k^2) f(u) + \dots \right]$$

In general the model becomes (with $\bar{h}_i(\mu) = h_i(-\mu)$)


$$-\mathcal{S}_{\text{eff}} = \sum_i \lambda_i(u, \kappa, N_\tau) \mathcal{S}_i^S - 2N_f \sum_i \left[h_i(u, \kappa, \mu, N_\tau) \mathcal{S}_i^A + \bar{h}_i(u, \kappa, \mu, N_\tau) \mathcal{S}_i^{\dagger A} \right]$$

Now, keep only $\lambda_1 \mathcal{S}_1^S$ and $h_1 \mathcal{S}_1^A + \bar{h}_1 \mathcal{S}_1^{\dagger A}$ (now called just λ, h)

Higher powers of loops are resummed into a determinant:


$$\begin{aligned} Z_{\text{eff}}(\lambda_1, h_1, \bar{h}_1; N_\tau) &= \int [dL] \left(\prod_{\langle ij \rangle} [1 + 2\lambda_1 \text{Re} L_i L_j^*] \right) \\ &\quad \left(\prod_x \underbrace{\det[(1 + h_1 W_x)(1 + \bar{h}_1 W_x^\dagger)]^{2N_f}}_{\equiv Q(L_x, L_x^*)^{N_f}} \right) \end{aligned}$$

QCD: first order deconf. transition region




deconfinement p.t.:
 breaking of global $Z(3)$ symmetry;
 explicitly broken by quark masses
 transition weakens

Phase diagram in eff. theory:



Phase boundary in two-coupling space




$$\lambda_{pc}(h) = \lambda_0 - a_1 h$$



Observable to identify order of p.t.:

$$\delta B_Q = B_4(\delta Q) = \frac{\langle (\delta Q)^4 \rangle}{\langle (\delta Q)^2 \rangle^2}$$

$$B_4(x) = 1.604 + bL^{1/\nu}(x - x_c) + \dots$$



parameter along phase boundary




Critical point

$$\lambda_c = 0.18672(7), h_c = 0.000731(40)$$

Mapping back to QCD:

N_f	M_c/T	$\kappa_c(N_\tau = 4)$	$\kappa_c(4)$, Ref. [23]	$\kappa_c(4)$, Ref. [22]
1	7.22(5)	0.0822(11)	0.0783(4)	~ 0.08
2	7.91(5)	0.0691(9)	0.0658(3)	—
3	8.32(5)	0.0625(9)	0.0595(3)	—

Convergence properties:



Finite density: sign problem solvable

- Metropolis algorithm: Mild sign problem; $\frac{\mu}{T} \lesssim 3$
- Worm algorithm: No sign problem






Figure: Quark density calculated with Z_{eff} from Metropolis or worm algorithm on 24^3 lattices for $\frac{\mu}{T} = 1$ and 2.

Critical quark mass as function of chemical potential




- Conclusion: Deconfinement critical line (or surface if we allow different quark masses) determined for *all* chemical potentials

The fully calculated deconfinement transition



deconfinement critical surface

phase diagram for $N_f=2, N_t=6$




Outlook

- Heavy quarks delay baryon condensation , want physical quark masses
- Light quarks: need very large order hopping expansion, difficult
- Instead: compute corrections to the strong coupling limit of the chiral staggered quark action, U(1) chiral symmetry

$$S_F = \sum_{x,\nu=1,4} \eta_{x,\nu} \bar{\chi}_x \left[U_{x,\nu} \chi_{x+\nu} - U_{x-\nu,\nu}^\dagger \chi_{x-\nu} \right] + 2m_q \sum_x \bar{\chi}_x \chi_x$$

Links can be integrated, resulting monomer-dimer action simulated, worm algorithm



Liquid-gas transition
at infinite coupling, $m=0$
de Forcrand, Fromm 09

Conclusions

- Proposal for two-step treatment of QCD phase transition:
 - I. Derivation of effective action by strong-coupling expansion
 - II. Simulation of effective theory
- $Z(N)$ -invariant effective theory for Yang-Mills, correct order of p.t., crit. temperature $\sim 10\%$ accurate in the continuum limit
- Deconfinement for heavy fermions and all chemical potentials
- Hope for finite density QCD: Treatment of light fermions?