

Dilepton Production at SPS Energies

Hendrik van Hees

Texas A&M University

December 17, 2007



Outline

- 1 Introduction
- 2 Thermal Fireball
- 3 Sources of Dileptons
 - In-medium vector mesons (thermal source)
 - $q\bar{q}$ annihilation in the QGP (thermal source)
 - Multi-pion processes (thermal source)
 - Meson t-channel exchange (thermal source)
 - ρ decay after thermal freezeout
 - “Primordial” ρ mesons
 - Drell-Yan Annihilation and correlated charm decays
- 4 Comparison to NA60 data
 - Invariant-mass spectra
 - m_T spectra
 - Sensitivity to T_c and hadro-chemistry
 - Inverse-slope analysis
- 5 Conclusions and Outlook
- 6 Backup Slides

Electromagnetic probes in heavy-ion collisions

- γ, ℓ^\pm : no strong interactions
- reflect whole “history” of collision:
 - from pre-equilibrium phase
 - from thermalized medium
QGP and hot hadron gas
 - from VM decays after thermal freezeout







Fig. by A. Drees

In-medium spectral functions and baryon effects



- **baryon effects** important $\leftrightarrow N_B + N_{\bar{B}}$ relevant (not $N_B - N_{\bar{B}}$)
 - large contribution to broadening of the peak
 - responsible for most of the strength at small M

Fireball and Thermodynamics

- cylindrical fireball model: $V_{\text{FB}} = \pi(z_0 + v_{z0}t + \frac{a_z}{2}t^2) (\frac{a_{\perp}}{2}t^2 + r_0)^2$

- thermodynamics:

- isentropic expansion; S_{tot} fixed by N_{ch} ; $T_c = T_{\text{chem}} = 175$ MeV
- $T > T_c$: massless gas for QGP with $N_f^{\text{eff}} = 2.3$
- mixed phase: $f_{\text{HG}}(t) = [s_c^{\text{QGP}} - s(t)]/[s_c^{\text{QGP}} - s_c^{\text{HG}}]$
- $T < T_c$: hadron-resonance gas


- $\Rightarrow T(t), \mu_{\text{baryon,meson}}(t)$

- chemical freezeout:

- $\mu_N^{\text{chem}} = 232$ MeV
- hadron ratios fixed
 $\Rightarrow \mu_N, \mu_{\pi}, \mu_K, \mu_{\eta}$ at fixed
 $s/\rho_B = 27$

- thermal freezeout:

$$(T_{\text{fo}}, \mu_{\pi}^{\text{fo}}) \simeq (120, 80) \text{ MeV}$$



Flow and particle/resonance distributions

- assume **local thermal equilibrium**: $T(t)$
- collective **radial flow**: $u(t, \vec{x}) = 1/\sqrt{1 - \vec{v}^2}(1, \vec{v})$
- $\vec{v}(t, \vec{x}) = a_{\perp} t \vec{x}_{\perp}/R(t)$
- phase-space distribution for hadrons

$$\frac{dN_i}{d^3\vec{p}d^3\vec{x}} = \frac{g_i}{(2\pi)^3} f_{B/F} \left(\frac{p \cdot u(t, \vec{x})}{T(t)} \right) \exp \left(\frac{\mu_i(t)}{T(t)} \right)$$


- NB:
 - covariant notation $d^3\vec{x}d^3\vec{p} = p_{\mu}d\sigma^{\mu}d^3\vec{p}/\sqrt{\vec{p}^2 + m^2}$
 - $pu(t, \vec{x}) = \bar{p}_0$: energy of particle in **rest frame of fluid cell**
- phase-space distribution for **bosonic resonances**:

$$\frac{dN_i}{d^4pd^3\vec{x}} = \frac{g_i}{(2\pi)^4} f_B \left(\frac{p \cdot u(t, \vec{x})}{T(t)} \right) \exp \left(\frac{\mu_i(t)}{T(t)} \right) [-2p_0 \text{Im } D_i(p)]$$

- $D_i(p)$: propagator of resonance,
 $A_i(p) = -2 \text{Im } D_i(p)$: spectral function

Radiation from thermal sources: ρ decays

- model assumption: **vector-meson dominance**



$$\begin{aligned} \frac{dN_{\rho \rightarrow l+l^-}^{(\text{MT})}}{d^4x d^4q} &= \frac{M}{q^0} \Gamma_{\rho \rightarrow l+l^-}(M) \frac{dN_{\rho}}{d^3\vec{x} d^4q} \\ &= -\frac{\alpha^2}{3\pi^3} \frac{L(M^2)}{M^2} \frac{m_{\rho}^4}{g_{\rho}^2} g_{\mu\nu} \text{Im} D_{\rho}^{\mu\nu}(M, \vec{q}) f_B \left(\frac{q \cdot u}{T(t)} \right) \exp \left(\frac{2\mu_{\pi}(t)}{T(t)} \right) \end{aligned}$$

- special case of McLerran-Toimela (MT) formula
- $M^2 = q^2$: invariant mass, M , of dilepton pair
- $L(M^2) = (1 + 2m_l^2/M^2) \sqrt{1 - 4m_l^2/M^2}$: dilepton phase-space factor
- $D_{\rho}^{\mu\nu}(M, \vec{q})$: (four-transverse part of) in-medium ρ propagator at given $T(t)$, $\mu_{\text{meson/baryon}}(t)$
- analogous for ω and ϕ

Radiation from thermal sources: $q\bar{q}$ annihilation

- General: McLerran-Toimela formula

$$\frac{dN_{l+l-}^{(\text{MT})}}{d^4x d^4q} = -\frac{\alpha^2}{3\pi^3} \frac{L(M^2)}{M^2} g_{\mu\nu} \text{Im} \sum_i \Pi_{\text{em},i}^{\mu\nu}(M, \vec{q}) f_B \left(\frac{q \cdot u}{T(t)} \right) \exp \left(\frac{\mu_i(t)}{T(t)} \right)$$

- i enumerates partonic/hadronic sources of em. currents
- in-medium em. current-current correlation function

$$\Pi_{\text{em},i}^{\mu\nu} = i \int d^4x \exp(iqx) \Theta(x^0) \left\langle \left[j_{\text{em},i}^\mu(x), j_{\text{em},i}^\nu(x) \right]_- \right\rangle$$

- in QGP phase: $q\bar{q}$ annihilation
- HTL improved electromagnetic current correlator



$$-i\Pi_{\text{em},\text{QGP}} = \text{Diagram}$$

Radiation from thermal sources: multi- π processes

- use vector/axial-vector correlators from τ -decay data
- Dey-Elefsky-Ioffe mixing: $\hat{\varepsilon} = 1/2\varepsilon(T, \mu_\pi)/\varepsilon(T_c, 0)$


$$\Pi_V = (1 - \hat{\varepsilon})z_\pi^4 \Pi_{V,4\pi}^{\text{vac}} + \frac{\hat{\varepsilon}}{2}z_\pi^3 \Pi_{A,3\pi}^{\text{vac}} + \frac{\hat{\varepsilon}}{2}(z_\pi^4 + z_\pi^5)\Pi_{A,5\pi}^{\text{vac}}$$

- avoid double counting: leave out two-pion piece and $a_1 \rightarrow \rho + \pi$ (already contained in ρ spectral function)



Radiation from thermal sources: Meson t-channel exchange



- motivation: q_T spectra too soft compared to NA60 data
- **thermal contributions** not included in models so far



- also for π , a_1

Radiation from thermal sources: Meson t-channel exchange

- t-channel exchange contributions become significant at **high momenta**



ρ decay after thermal freezeout

- assume “sudden freezeout” at constant “lab time”: $t = t_{\text{fo}}$
- then Cooper-Frye formula with $d\sigma^\mu = (d^3\vec{x}, 0, 0, 0)$

$$\begin{aligned}\frac{dN_{\rho \rightarrow l^+l^-}^{(\text{fo})}}{d^3\vec{x}d^4\vec{q}} &= \frac{\Gamma_{l^+l^-}}{\Gamma_\rho^{\text{tot}}} \frac{dN_i}{d^3\vec{x}d^4q} \\ &= \frac{q_0}{M} \frac{1}{\Gamma_\rho^{\text{tot}}} \left[\frac{dN_{\rho \rightarrow l^+l^-}^{(\text{MT})}}{d^4xd^4q} \right]_{t=t_{\text{fo}}}\end{aligned}$$

- use vacuum ρ shape with in-medium width $\Gamma_\rho^{\text{tot}} \simeq 260 \text{ MeV}$
- NB: Momentum dependence for dilepton spectra from ρ decays after thermal freezeout:

like hadron spectra!


- $\Leftrightarrow l^+l^-$ from thermal sources softer by Lorentz factor M/q^0 compared to l^+l^- from decay of freeze-out ρ 's

Decay of “primordial” ρ mesons

- ρ mesons, escaping from the fireball **without thermalization**
- pp data for **initial ρ spectra**; **Cronin effect** via “Gaussian smearing”
- Schematic **jet-quenching model**

$$P_{\text{esc}} = \exp \left(- \int dt \sigma_{\rho}^{\text{abs}}(t) \varrho(t) \right),$$

$$\sigma_{\rho}^{\text{abs}}(t) = \begin{cases} \sigma_{\text{ph}} = 0.4 \text{ mb} & \text{for } t < q_0/m_{\rho} \tau_{\text{f}} \\ \sigma_{\text{had}} = 5 \text{ mb} & \text{for } t > q_0/m_{\rho} \tau_{\text{f}} \end{cases}$$



- check with **pion R_{AA}** data
- “primordial ρ ’s” + freezeout ρ ’s
- **hard q_T spectra**
including jet quenching

- **invariant-mass spectrum** for DY pairs

$$\left. \frac{dN_{\text{DY}}^{AA}}{dMdy} \right|_{b=0} = \frac{3}{4\pi R_0^2} A^{4/3} \frac{d\sigma_{\text{DY}}^{NN}}{dMdy}$$
$$\frac{d\sigma_{\text{DY}}^{NN}}{dMdy} = K \frac{8\pi\alpha}{9sM} \sum_{q=u,d,s} e_q^2 [q(x_1)\bar{q}(x_2) + \bar{q}(x_1)q(x_2)]$$

- **parton distribution functions**: GRV94LO
- **higher-order effects**
 - K factor
 - non-zero pair q_T : for IMR and HMR fitted by **Gaussian spectrum** (NA50 procedure)
- extrapolation to LMR: constrained by photon point $M \rightarrow 0$



- **invariant-mass spectrum** for DY pairs

$$\left. \frac{dN_{\text{DY}}^{AA}}{dMdy} \right|_{b=0} = \frac{3}{4\pi R_0^2} A^{4/3} \frac{d\sigma_{\text{DY}}^{NN}}{dMdy}$$
$$\frac{d\sigma_{\text{DY}}^{NN}}{dMdy} = K \frac{8\pi\alpha}{9sM} \sum_{q=u,d,s} e_q^2 [q(x_1)\bar{q}(x_2) + \bar{q}(x_1)q(x_2)]$$

- **parton distribution functions**: GRV94LO
- **higher-order effects**
 - K factor
 - non-zero pair q_T : for IMR and HMR fitted by **Gaussian spectrum** (NA50 procedure)
- extrapolation to LMR: constrained by photon point $M \rightarrow 0$
- Correlated decays of D and \bar{D} mesons
 - use data (provided by NA60 collaboration)


Invariant-mass spectra

- Fireball with “standard” EoS-A ($T_c = T_{\text{chem}} = 175$ MeV)
- overall normalization \Leftrightarrow total fireball lifetime
- relative normalization of thermal radiation fixed by rates
- rates integrated over time, volume, \vec{q} including NA60 acceptance




- good description of data

Excess spectra: IMR and multi-pion contributions



- “ 4π contributions” ($\pi + \omega, a_1 \rightarrow \mu^+ + \mu^-$)
- slightly enhanced by VA mixing





Excess spectra: baryon effects






- in-medium VM spectral functions without baryon effects

- not enough broadening around $M = m_{\rho}^{\text{vac}}$
- lack of strength at low mass $M \rightarrow m_{\text{thr}} = 2m_{\mu}$

Excess spectra: q_T binning






m_T spectra (central)






- fixed normalization in $0 \leq q_T \leq 0.5$ GeV bin
- satisfactory description of data
- high-mass bin slightly overestimated
- hard probes important for $q_T \geq 1$ GeV

m_T spectra (semicentral)



- theoretical spectra too soft in low-mass and ρ -region bin
- room for more “primordial ρ ’s”



Sensitivity to meson t -channel exchange contributions



- use $2\times$ of ω - t exchange to account for other mesons (e.g., a_1 , π)
- **hardest among thermal sources**
- absolute strength not sufficient to resolve discrepancy with data

Sensitivity to T_c and hadro-chemistry



- recent lattice QCD: $T_c \simeq 190\text{-}200$ MeV or $T_c \simeq 150\text{-}160$ MeV?
- thermal-model fits to hadron ratios: $T_{\text{chem}} \simeq 150\text{-}160$ MeV




- EoS-A: $T_c = T_{\text{chem}} = 175$ MeV
- EoS-B: $T_c = T_{\text{chem}} = 160$ MeV
- EoS-C: $T_c = 190$ MeV, $T_{\text{chem}} = 160$ MeV
 - $T_c \geq T \geq T_{\text{chem}}$: hadron gas in chemical equilibrium
- keep fireball parameters the same (including life time)

Sensitivity to T_c and hadro-chemistry


- recent lattice QCD: $T_c \simeq 190\text{-}200$ MeV or $T_c \simeq 150\text{-}160$ MeV?
- thermal-model fits to hadron ratios: $T_{\text{chem}} \simeq 150\text{-}160$ MeV



- EoS-A: $T_c = T_{\text{chem}} = 175$ MeV
- EoS-B: $T_c = T_{\text{chem}} = 160$ MeV
- EoS-C: $T_c = 190$ MeV, $T_{\text{chem}} = 160$ MeV
 - $T_c \geq T \geq T_{\text{chem}}$: hadron gas in chemical equilibrium
- keep fireball parameters the same (including life time)





- mass spectra comparable to EoS-A \leftrightarrow slight enhancement of fireball lifetime
- in IMR **QGP** > multi-pion contribution
- higher hadronic temperatures \Rightarrow slightly harder q_T spectra
- not enough to resolve discrepancy with data



- mass spectra comparable to EoS-A \leftrightarrow slight reduction of fireball lifetime
- in IMR **multi-pion** \gg **QGP** contribution
- higher hadronic temperatures + high-density hadronic phase \Rightarrow harder q_T spectra
- better agreement with data

IMR: QGP vs. multi-pion radiation



- **EoS-B**: QGP dominates over multi-pion radiation
- opposite in **EoS-A** and **EoS-C**
- multi-pion radiation **dominantly from high-density hadronic phase**

reason: $dN_{ll}/dMdT \propto \text{Im} \Pi_{\text{em}}(M, T) \exp(-M/T) T^{-5.5}$



- radiation maximal for $T = T_{\text{max}} = M/5.5$
 - hadronic and partonic radiation “dual” for $T \sim T_c$
- compatible with chiral-symmetry restoration!**

Inverse-slope analysis

- to extract T_{eff} fit to



$$\frac{1}{q_T} \frac{dN}{dq_T} = \frac{1}{m_T} \frac{dN}{dm_T} = C \exp\left(-\frac{m_T}{T_{\text{eff}}}\right)$$

- fit of theoretical q_T spectra: $1 \text{ GeV} < q_T < 1.8 \text{ GeV}$





- standard fireball acceleration: **too soft q_T spectra**
- lower T_c in EoS-B and EoS-C helps (higher hadronic temperatures)
- NB: here, Drell Yan contribution taken out

Inverse-slope analysis



- enhance fireball acceleration to $a_{\perp} = 0.1c^2/\text{fm}$
- effective at all stages of fireball evolution
- agreement in IMR not spoiled \Leftrightarrow dominated from earlier stages
- EoS-B harder \Leftrightarrow relative contribution of harder freezeout ρ decays vs. thermal ρ 's larger


Inverse-slope analysis



- sensitivity to contributions from meson t -channel exchange
 - hardens low-mass region
 - using vacuum ρ in t -channel contribution: enhances slope in ρ region
- sensitivity to Drell-Yan contribution
 - for IMR: describes effect seen in data (open vs. solid square data point)
 - in LMR: too high around muon threshold \Leftrightarrow due to uncertainties in extrapolation to low M ?!?

Slopes of thermal radiation


- use EoS-B with $a_{\perp} = 0.1c^2/\text{fm}$
- fit the dilepton slopes from thermal sources only



- freeze-out and primordial ρ 's crucial for $q_T > 1$ GeV

Hadron slopes

- hadron slopes extracted by fit to **freeze-out contribution**



- η , ω , and ϕ slopes described by EoS-A and **smaller a_{\perp}**
- ρ slope by EoS-B with **larger a_{\perp}**
- possible resolution: **“successive freezeout”** \Leftrightarrow vector mesons freeze out at different times, determined by interaction strength (i.e., width)

Conclusions and Outlook


- dilepton spectra \Leftrightarrow in-medium em. current correlator
- model for dilepton sources
 - radiation from thermal sources: QGP, ρ , ω , ϕ
 - ρ -decay after thermal freeze-out
 - decays of non-thermalized primordial ρ 's
 - Drell-Yan annihilation, correlated $D\bar{D}$ decays

Conclusions and Outlook

- dilepton spectra \Leftrightarrow in-medium em. current correlator
- model for dilepton sources
 - radiation from thermal sources: QGP, ρ , ω , ϕ
 - ρ -decay after thermal freeze-out
 - decays of non-thermalized primordial ρ 's
 - Drell-Yan annihilation, correlated $D\bar{D}$ decays
- invariant-mass spectra and medium effects
 - excess yield dominated by radiation from thermal sources
 - baryons essential for in-medium properties of vector mesons
 - melting ρ with little mass shift robust signal! (independent of T_c)
 - IMR well described by scenarios with radiation dominated either by QGP or multi-pion processes (depending on EoS)
 - Reason: mostly from thermal radiation around $160 \text{ MeV} \leq T \leq 190 \text{ MeV}$
 - \Leftrightarrow "parton-hadron" duality of rates
 - \Leftrightarrow compatible with chiral-symmetry restoration!
 - dimuons in In-In (NA60), Pb-Au (CERES/NA45), γ in Pb-Pb (WA98)

- fireball/freeze-out dynamics $\Leftrightarrow m_T$ spectra and effective slopes
 - “non-thermal sources” important for $q_T \gtrsim 1$ GeV
 - lower $T_c \Rightarrow$ higher hadronic temperatures \Rightarrow harder q_T spectra
 - to describe measured effective slopes $a_{\perp} = 0.085c^2/\text{fm} \rightarrow 0.1c^2/\text{fm}$
 - off-equilibrium effects (viscous hydro)?
- Further developments
 - understand recent PHENIX results (large dilepton excess in LMR)
 - **vector**- should be complemented with **axial-vector**-spectral functions (a_1 as chiral partner of ρ)
 - constrained with IQCD via **in-medium Weinberg chiral sum rules**
 - **direct connection to chiral phase transition!**



Backup: parton-hadron duality of rates



- in-medium **hadron gas** matches with **QGP**
- similar results also for γ rates
- “quark-hadron duality”!?
- indirect evidence for **chiral-symmetry restoration**

Backup: CERES/NA45 dielectron spectra

- good agreement also for **dielectron** spectra in 158 GeV Pb-Au
- allows further check of **low-mass tail from baryon effects** down to $M \rightarrow 2m_e$



Backup: Low- m_T rise

- observed low- q_T deviation from $\frac{dN}{m_T dm_T} \propto \exp\left(-\frac{m_T}{T_{\text{eff}}}\right)$
- asymptotic limit for $M \gg T$ AND $q_T \gg M$ for ll from fo- ρ decays
- for hadrons or radiation from freeze-out ρ 's:

$$\frac{dN(\text{fo})}{m_T dm_T} \propto \begin{cases} \exp\left(-\frac{m_T}{T_{\text{eff}}}\right) & \text{for } q_T \gg M \gg T \\ \sqrt{m_T} \exp\left(-\frac{m_T}{T'_{\text{eff}}}\right) & \text{for } q_T \ll T \ll M. \end{cases}$$

- for radiation from thermal source:

$$\frac{dN(\text{MT})}{m_T dm_T} \propto \begin{cases} \frac{1}{m_T} \exp\left(-\frac{m_T}{T_{\text{eff}}}\right) & \text{for } q_T \gg M \gg T \\ \frac{1}{\sqrt{m_T}} \exp\left(-\frac{m_T}{T'_{\text{eff}}}\right) & \text{for } q_T \ll T \ll M. \end{cases}$$

- with effective inverse slopes

$$T_{\text{eff}} = T \sqrt{\frac{1 + \xi \beta^B}{1 - \xi \beta^B}} \quad (0 < \xi < 1), \quad T'_{\text{eff}} \simeq T + \frac{M}{2} \langle \beta^2(r) \rangle_r = T + \frac{M}{4} \beta_B^2.$$

- possibly also effect of Bose Enhancement due to μ_{mesons}

Backup: Low- m_T rise

